

化学療法レジメンセット登録用紙

領域	血液腫瘍	ルート	リザーバー
適応	【AML】急性骨髄性白血病		
No(3桁)	019	CODE	004002001019
診療科	内科	医師	堤
セット名称	AraC+DNR+6MP		
申請年月日	平成21年7月7日		
緊急		通常	●

抗癌剤名称	手技	標準投与量
シタラビン	CIV	80 mg/m ²
ダウノマイシン	DIV	40 mg/m ²
ロイケリン	p.o	70 mg/m ²

Aポート	Vポート	末梢	確認		
	●		医師	看護師	薬剤師

投与順	Rp	投与時間	投与スケジュール: (投 休) or (Interval日数 日) 《 》 クール																												
			-1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28
1	ロイケリン散()mg 手技: 経路:朝・夕食後	分2 p.o		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
2	グラニセロンパック 50mL 手技: 静脈内 経路:	30分 DIV		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
3	ダウノマイシン()mg + 生食100ml 手技: 静脈内 経路:	30分 DIV		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
4	シタラビン()mg + 生食500ml 手技: 静脈内 経路:	24時間 CIV		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
5	ソルデム3A 1000ml 手技: 静脈内 経路:	24時間 CIV		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
6	ヘパリンNa 1筒 手技: 静脈内 経路:シタラビン点滴終了後に	IV		●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
7	手技: 経路:			●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
8	手技: 経路:			●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
9	手技: 経路:			●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●
10	手技: 経路:			●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●	●