## 化学療法レジメンセット登録用紙

領域	血液腫瘍	ルート	リザーバー								
適応	【NHL】非	リンパ腫									
No(3桁)	007	CODE	004012001007								
診療科	内科	医師	堤								
セット名称											
申請年月日	平瓦	平成21年7月7日									
緊急		通常	•								

抗癌剤名称	手技	標準	<b>準投</b> 与量
オンコビン	DIV	1.4	mg/m <sup>*</sup>
ドキソルビシン	DIV	50	mg/ <b>m</b> i
エンドキサン	DIV	750	mg/m <sup>²</sup>
プレドニン	p.o	100	mg/body

Aポート Vポー	<b>−</b> ト  ;	末梢		確認									
			医師	看護師	薬剤師								
	'												
	•												

			7	投与.	スク	ァジニ	1-	ル:	:	(		;	投			休	)	or	(Iı	nter	val	日数	数			日)	<b>«</b>	<u> </u>	<u></u>	7—J	7
投与順	Rp	投与時間	-1	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23 2	24 2	25 2	6 2	27 28	3
1	プレドニン錠20錠 毎食後(朝 錠-昼 錠-タ 釘			•																									T		1
	手技: 経路:	p.o																													
2	グラニセトロンバック 50mL	30分																													
	手技:静脈内 経路:	DIV																													
3	生食50mL+オンコビン( )mg	15分		•																											1
١	手技:静脈内 経路:	DIV																													
4	生食100mL+ドキソルビシン( )mg	1時間		•																											1
4	手技:静脈内 経路:	DIV																													
5	生食250mL+エンドキサン( )mg	3時間																													1
	手技:静脈内 経路:溶解後6時間以内に投与終了する事	DIV																													
6	ソルデム3A 1000mL	6時間		•																											1
"	手技:静脈内 経路:	DIV								- 1																					
7	ヘパリンNaロックシリンジ 1本																												T		1
_ ′	手技:静脈内 経路:	IV																													
8																													T		1
l °	手技: 経路:																														
9																															1
9	手技: 経路:																												$\perp$		
10								Ī																							
	手技: 経路:																														